

# EXEMPLAR SOLUTIONS MATHS

Chapter 5 : Continuity and Differentiability

Class  
**12**



Powered by



## Chapter 5: Continuity and Differentiability

### Exercise 5.3

**Short Answer (S.A.)**

**1. Examine the continuity of the function  $f(x) = x^3 + 2x^2 - 1$  at  $x = 1$**

**Solution:**

We know that,  $y = f(x)$  will be continuous at  $x = a$  if,

$$\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a^+} f(x)$$

Given:  $f(x) = x^3 + 2x^2 - 1$

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{h \rightarrow 0} (1+h)^3 + 2(1+h)^2 - 1 = 1 + 2 - 1 = 2$$

$$\lim_{x \rightarrow 1} f(x) = (1)^3 + 2(1)^2 - 1 = 1 + 2 - 1 = 2$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{h \rightarrow 0} (1+h)^3 + 2(1+h)^2 - 1 = 1 + 2 - 1 = 2$$

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1^+} f(x) = 2.$$

Thus,  $f(x)$  is continuous at  $x = 1$ .

**Find which of the functions in Exercises 2 to 10 is continuous or discontinuous at the indicated points:**

2.  $f(x) = \begin{cases} 3x+5, & \text{if } x \geq 2 \\ x^2, & \text{if } x < 2 \end{cases}$  at  $x = 2$

**Solution:**

Checking the continuity of the given function, we have

$$\lim_{x \rightarrow 2^+} f(x) = 3x + 5 = \lim_{h \rightarrow 0} 3(2+h) + 5 = 11$$

$$\lim_{x \rightarrow 2} f(x) = 3x + 5 = 3(2) + 5 = 11$$

$$\lim_{x \rightarrow 2^-} f(x) = x^2 = \lim_{h \rightarrow 0} (2-h)^2 = \lim_{h \rightarrow 0} (2)^2 + h^2 - 4h = (2)^2 = 4$$

Now, since  $\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2} f(x) \neq \lim_{x \rightarrow 2^+} f(x)$

Thus,  $f(x)$  is discontinuous at  $x = 2$ .

3.  $f(x) = \begin{cases} \frac{1 - \cos 2x}{x^2}, & \text{if } x \neq 0 \\ 5, & \text{if } x = 0 \end{cases}$  at  $x = 0$

**Solution:**



Checking the right hand and left hand limits of the given function, we have

$$\begin{aligned}\lim_{x \rightarrow 0^-} f(x) &= \frac{1 - \cos 2x}{x^2} \\&= \lim_{h \rightarrow 0} \frac{1 - \cos 2(0 - h)}{(0 - h)^2} = \lim_{h \rightarrow 0} \frac{1 - \cos(-2h)}{h^2} \\&= \lim_{h \rightarrow 0} \frac{1 - \cos 2h}{h^2} \\&= \lim_{h \rightarrow 0} \frac{2 \sin^2 h}{h^2} \quad \left[ \because 1 - \cos \theta = 2 \sin^2 \frac{\theta}{2} \right] \\&= \lim_{h \rightarrow 0} \frac{2 \sin h}{h} \cdot \frac{\sin h}{h} = 2.1.1 = 2 \quad \left[ \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right]\end{aligned}$$

$$\begin{aligned}\lim_{x \rightarrow 0^+} f(x) &= \frac{1 - \cos 2x}{x^2} \\&= \lim_{h \rightarrow 0} \frac{1 - \cos 2(0 + h)}{(0 + h)^2} = \lim_{h \rightarrow 0} \frac{1 - \cos 2h}{h^2} \\&= \lim_{h \rightarrow 0} \frac{2 \sin^2 h}{h^2} = \frac{2 \sin h}{h} \cdot \frac{\sin h}{h} = 2.1.1 = 2\end{aligned}$$

$$\lim_{x \rightarrow 0} f(x) = 5$$

$$\text{As } \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} f(x) \neq \lim_{x \rightarrow 0^+} f(x)$$

Therefore, the given function  $f(x)$  is discontinuous at  $x = 0$ .

$$f(x) = \begin{cases} \frac{2x^2 - 3x - 2}{x - 2}, & \text{if } x \neq 2 \\ 5, & \text{if } x = 2 \end{cases} \quad \text{at } x = 2$$

4.

**Solution:**

The given function at  $x \neq 0$  can be rewritten as,

$$\begin{aligned}f(x) &= \frac{2x^2 - 3x - 2}{x - 2} \\&= \frac{2x^2 - 4x + x - 2}{x - 2} = \frac{2x(x - 2) + 1(x - 2)}{x - 2} \\&= \frac{(2x + 1)(x - 2)}{x - 2} = 2x + 1\end{aligned}$$

Now,

$$\begin{aligned}\lim_{x \rightarrow 2^-} f(x) &= 2x + 1 \\&= \lim_{h \rightarrow 0} 2(2 - h) + 1 = 4 + 1 = 5\end{aligned}$$

$$\begin{aligned}\lim_{x \rightarrow 2^+} f(x) &= 2x + 1 \\&= \lim_{h \rightarrow 0} 2(2 + h) + 1 = 4 + 1 = 5\end{aligned}$$

$$\lim_{x \rightarrow 2} f(x) = 5$$

$$\text{As } \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2} f(x) = 5$$

Thus,  $f(x)$  is continuous at  $x = 2$ .

$$f(x) = \begin{cases} \frac{|x-4|}{2(x-4)}, & \text{if } x \neq 4 \\ 0, & \text{if } x = 4 \end{cases} \text{ at } x = 4$$

5.

**Solution:**

Checking the right hand and left hand limits for the given function, we have

$$\begin{aligned} \lim_{x \rightarrow 4^-} f(x) &= \frac{|x-4|}{2(x-4)} \quad \left[ \begin{array}{l} \text{for } x < 4, |x-4| = -(x-4) \\ \text{for } x > 4, |x-4| = (x-4) \end{array} \right] \\ &= \lim_{h \rightarrow 0} \frac{-(4-h-4)}{2[4-h-4]} = \lim_{h \rightarrow 0} \frac{h}{-2h} = -\frac{1}{2} \end{aligned}$$

$$\lim_{x \rightarrow 4^+} f(x) = \frac{|x-4|}{2(x-4)} = \lim_{h \rightarrow 0} \frac{[4+h-4]}{2[4+h-4]} = \frac{1}{2}$$

$$\lim_{x \rightarrow 4} f(x) = 0$$

$$\therefore \lim_{x \rightarrow 4^-} f(x) \neq \lim_{x \rightarrow 4^+} f(x) \neq \lim_{x \rightarrow 4} f(x)$$

Thus,  $f(x)$  is discontinuous at  $x = 4$ .

$$f(x) = \begin{cases} |x| \cos \frac{1}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases} \text{ at } x = 0$$

6.

**Solution:**

Checking the right hand and left hand limits for the given function, we have

$$\begin{aligned} \lim_{x \rightarrow 0^-} f(x) &= |x| \cos \frac{1}{x} \\ &= \lim_{h \rightarrow 0} |0-h| \cos \frac{1}{(0-h)} = \lim_{h \rightarrow 0} h \cos \frac{1}{h} \\ &= 0 \quad \left[ \because \cos \frac{1}{x} \text{ oscillate between } -1 \text{ and } 1 \right] \end{aligned}$$

$$\lim_{x \rightarrow 0^+} f(x) = |x| \cos \frac{1}{x}$$

$$= \lim_{h \rightarrow 0} |0 + h| \cos \frac{1}{(0 + h)} = \lim_{h \rightarrow 0} h \cdot \cos \frac{1}{h} = 0$$

$$\lim_{x \rightarrow 0^+} f(x) = 0$$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0} f(x) = 0$$

Thus, the given function  $f(x)$  is continuous at  $x = 0$ .

$$f(x) = \begin{cases} |x-a| \sin \frac{1}{x-a}, & \text{if } x \neq a \\ 0, & \text{if } x = a \end{cases} \quad \text{at } x = a$$

7.

**Solution:**

Checking the right hand and left hand limits for the given function, we have

$$\begin{aligned} \lim_{x \rightarrow a^-} f(x) &= |x-a| \sin \frac{1}{x-a} \\ &= \lim_{h \rightarrow 0} |a-h-a| \cdot \sin \frac{1}{a-h-a} = \lim_{h \rightarrow 0} h \cdot \sin \frac{1}{-h} \\ &= \lim_{h \rightarrow 0} -h \cdot \sin \frac{1}{h} \quad [\because \sin(-\theta) = -\sin \theta] \\ &= 0 \times [\text{a number oscillating between } -1 \text{ and } 1] \\ &= 0 \end{aligned}$$

$$\begin{aligned} \lim_{x \rightarrow a^+} f(x) &= |x-a| \sin \frac{1}{x-a} \\ &= \lim_{h \rightarrow 0} |a+h-a| \cdot \sin \frac{1}{a+h-a} = \lim_{h \rightarrow 0} h \cdot \sin \frac{1}{h} \\ &= 0 \times [\text{a number oscillating between } -1 \text{ and } 1] \end{aligned}$$

$$\lim_{x \rightarrow a} f(x) = 0$$

Now, as

$$\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow a} f(x) = 0$$

Thus, the given function  $f(x)$  is continuous at  $x = 0$ .

$$f(x) = \begin{cases} \frac{e^x}{1+e^x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases} \quad \text{at } x = 0$$

8.

**Solution:**

Checking the right hand and left hand limits for the given function, we have

$$\begin{aligned}
 \lim_{x \rightarrow 0} f(x) &= \frac{e^{1/x}}{1 + e^{1/x}} \\
 &= \lim_{h \rightarrow 0} \frac{e^{\frac{1}{0-h}}}{1 + e^{\frac{1}{0-h}}} = \lim_{h \rightarrow 0} \frac{e^{-1/h}}{1 + e^{-1/h}} \\
 &= \lim_{h \rightarrow 0} \frac{1}{e^{1/h} (1 + e^{-1/h})} = \lim_{h \rightarrow 0} \frac{1}{e^{1/h} - 1} = \frac{1}{e^{1/0} - 1} \\
 &= \frac{1}{e^{\infty} - 1} = \frac{1}{0 - 1} = -1 \quad [\because e^{\infty} = 0]
 \end{aligned}$$

$$\begin{aligned}
 \lim_{x \rightarrow 0^+} f(x) &= \frac{e^{1/x}}{1 + e^{1/x}} \\
 &= \lim_{h \rightarrow 0} \frac{e^{\frac{1}{0+h}}}{1 + e^{\frac{1}{0+h}}} = \lim_{h \rightarrow 0} \frac{e^{1/h}}{1 + e^{1/h}} \\
 &= \lim_{h \rightarrow 0} \frac{1}{e^{-1/h} (1 + e^{1/h})} = \lim_{h \rightarrow 0} \frac{1}{e^{-1/h} + 1} \\
 &= \frac{1}{e^{-\infty} + 1} = \frac{1}{0 + 1} = 1 \quad [e^{-\infty} = 0]
 \end{aligned}$$

$$\lim_{x \rightarrow 0} f(x) = 0$$

Now, as

$$\lim_{x \rightarrow 0^+} f(x) \neq \lim_{x \rightarrow 0} f(x) \neq \lim_{x \rightarrow 0^-} f(x)$$

Thus,  $f(x)$  is discontinuous at  $x = 0$ .

$$f(x) = \begin{cases} \frac{x^2}{2}, & \text{if } 0 \leq x \leq 1 \\ 2x^2 - 3x + \frac{3}{2}, & \text{if } 1 < x \leq 2 \end{cases} \quad \text{at } x = 1$$

9.

**Solution:**

Checking the right hand and left hand limits for the given function, we have

$$\lim_{x \rightarrow 1^-} f(x) = \frac{x^2}{2} = \lim_{h \rightarrow 0} \frac{(1-h)^2}{2} = \frac{1}{2}$$

$$\lim_{x \rightarrow 1} f(x) = \frac{x^2}{2} = \frac{(1)^2}{2} = \frac{1}{2}$$



$$\lim_{x \rightarrow 1^-} f(x) = 2x^2 - 3x + \frac{3}{2} = 2(1)^2 - 3(1) + \frac{3}{2} = 2 - 3 + \frac{3}{2} = \frac{1}{2}$$

Now, as

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1} f(x) = \frac{1}{2}$$

Thus, the given function  $f(x)$  is continuous at  $x = 1$ .

**10.  $f(x) = |x| + |x - 1|$  at  $x = 1$**

**Solution:**

Checking the right hand and left hand limits for the given function, we have

$$\begin{aligned} \lim_{x \rightarrow 1^-} f(x) &= |x| + |x - 1| = \lim_{h \rightarrow 0} |1 - h| + |1 - h - 1| \\ &= |1 - 0| + |1 - 0 - 1| = 1 + 0 = 1 \end{aligned}$$

$$\begin{aligned} \lim_{x \rightarrow 1^+} f(x) &= |x| + |x - 1| \\ &= \lim_{h \rightarrow 0} |1 + h| + |1 + h - 1| = 1 + 0 = 1 \end{aligned}$$

$$\lim_{x \rightarrow 1} f(x) = |x| + |x - 1| = |1| + |1 - 1| = 1 + 0 = 1$$

Now, as

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1} f(x)$$

Thus,  $f(x)$  is continuous at  $x = 1$ .

**Find the value of  $k$  in each of the Exercises 11 to 14 so that the function  $f$  is continuous at the indicated point:**

$$f(x) = \begin{cases} 3x - 8, & \text{if } x \leq 5 \\ 2k, & \text{if } x > 5 \end{cases} \quad \text{at } x = 5$$

**11.**

**Solution:**

Finding the left hand and right hand limits for the given function, we have

$$\begin{aligned} \lim_{x \rightarrow 5^-} f(x) &= 3x - 8 \\ &= \lim_{h \rightarrow 0} 3(5 - h) - 8 = 15 - 8 = 7 \end{aligned}$$

$$\lim_{x \rightarrow 5^+} f(x) = 2k$$

As the function is continuous at  $x = 5$

$$\lim_{x \rightarrow 5^-} f(x) = \lim_{x \rightarrow 5^+} f(x)$$

So,

$$7 = 2k$$

$$k = 7/2 = 3.5$$

Therefore, the value of  $k$  is 3.5

$$f(x) = \begin{cases} \frac{2^{x+2} - 16}{4^x - 16}, & \text{if } x \neq 2 \\ k & , \text{ if } x = 2 \end{cases} \text{ at } x = 2$$

12.

**Solution:**

The given function  $f(x)$  can be rewritten as,

$$f(x) = \frac{2^{x+2} - 16}{4^x - 16} = \frac{2^2 \cdot 2^x - 16}{(2^x)^2 - (4)^2} = \frac{4(2^x - 4)}{(2^x - 4)(2^x + 4)}$$

$$f(x) = \frac{4}{2^x + 4}$$

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{h \rightarrow 0} \frac{4}{2^{2-h} + 4} = \frac{4}{2^2 + 4} = \frac{4}{4 + 4} = \frac{4}{8} = \frac{1}{2}$$

$$\lim_{x \rightarrow 2} f(x) = k$$

As the function is continuous at  $x = 2$ .

$$\therefore \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2} f(x)$$

So,  $k = \frac{1}{2}$

Therefore, the value of  $k$  is  $\frac{1}{2}$

$$f(x) = \begin{cases} \frac{\sqrt{1+kx} - \sqrt{1-kx}}{x}, & \text{if } -1 \leq x < 0 \\ \frac{2x+1}{x-1}, & \text{if } 0 \leq x \leq 1 \end{cases} \text{ at } x = 0$$

13.

**Solution:**

Finding the left hand and right hand limits for the given function, we have

$$\begin{aligned} \lim_{x \rightarrow 0^+} f(x) &= \frac{\sqrt{1+kx} - \sqrt{1-kx}}{x} \\ &= \lim_{x \rightarrow 0^+} \frac{\sqrt{1+kx} - \sqrt{1-kx}}{x} \times \frac{\sqrt{1+kx} + \sqrt{1-kx}}{\sqrt{1+kx} + \sqrt{1-kx}} \\ &= \lim_{x \rightarrow 0^+} \frac{(1+kx) - (1-kx)}{x [\sqrt{1+kx} + \sqrt{1-kx}]} \\ &= \lim_{x \rightarrow 0^+} \frac{1+kx - 1+kx}{x [\sqrt{1+kx} + \sqrt{1-kx}]} \\ &= \lim_{x \rightarrow 0^+} \frac{2kx}{x [\sqrt{1+kx} + \sqrt{1-kx}]} \end{aligned}$$

$$\begin{aligned}
 &= \lim_{x \rightarrow 0^+} \frac{2k}{\sqrt{1+kx} + \sqrt{1-kx}} \\
 &= \lim_{h \rightarrow 0} \frac{2k}{\sqrt{1+k(0-h)} + \sqrt{1-k(0-h)}} \\
 &= \frac{2k}{\sqrt{1} + \sqrt{1}} = \frac{2k}{2} = k
 \end{aligned}$$

$$\lim_{x \rightarrow 0^-} f(x) = \frac{2x+1}{x-1} = \frac{2(0)+1}{0-1} = \frac{1}{-1} = -1$$

As the function is continuous at  $x = 0$ ,

$$\therefore \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^-} f(x)$$

$$k = -1$$

Therefore, the value of  $k$  is  $-1$

$$f(x) = \begin{cases} \frac{1 - \cos kx}{x \sin x}, & \text{if } x \neq 0 \\ \frac{1}{2}, & \text{if } x = 0 \end{cases}$$

14.

**Solution:**

Finding the left hand and right hand limits for the given function, we have

$$\begin{aligned}
 \lim_{x \rightarrow 0^+} f(x) &= \frac{1 - \cos kx}{x \sin x} \\
 &= \lim_{h \rightarrow 0} \frac{1 - \cos k(0-h)}{(0-h) \sin(0-h)} = \lim_{h \rightarrow 0} \frac{1 - \cos(-kh)}{-h \sin(-h)} \\
 &= \lim_{h \rightarrow 0} \frac{1 - \cos kh}{h \sin h} \quad \left[ \begin{array}{l} \because \sin(-\theta) = -\sin \theta \\ \cos(-\theta) = \cos \theta \end{array} \right] \\
 &= \lim_{h \rightarrow 0} \frac{2 \sin^2 \frac{kh}{2}}{h \sin h} \\
 &= \lim_{\substack{h \rightarrow 0 \\ kh \rightarrow 0}} \frac{2 \sin \frac{kh}{2}}{\frac{kh}{2}} \times \frac{\frac{kh}{2}}{2} \times \frac{\sin \frac{kh}{2}}{\frac{kh}{2}} \times \frac{kh}{2} \times \frac{1}{h \cdot \frac{\sin h}{h} \cdot h} \\
 &= 2 \cdot 1 \cdot \frac{kh}{2} \cdot 1 \cdot \frac{kh}{2} \cdot \frac{1}{h^2} \cdot 1 \quad \left[ \begin{array}{l} \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1 \text{ and} \\ \lim_{kh \rightarrow 0} \frac{\sin kh}{kh} = 1 \end{array} \right] \\
 &= \frac{k^2}{2}
 \end{aligned}$$

$$\lim_{x \rightarrow 0^-} f(x) = \frac{1}{2}$$

But, as the function is continuous we have

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x)$$

$$\text{So, } \frac{k^2}{2} = \frac{1}{2}$$

$$k^2 = 1 \Rightarrow k = \pm 1$$

Therefore, the value of  $k$  is  $\pm 1$

**15. Prove that the function  $f$  defined by**

$$f(x) = \begin{cases} \frac{x}{|x| + 2x^2}, & x \neq 0 \\ k, & x = 0 \end{cases}$$

**remains discontinuous at  $x = 0$ , regardless the choice of  $k$ .**

**Solution:**

Finding the left hand and right hand limit for the given function, we have

$$\begin{aligned} \lim_{x \rightarrow 0^-} f(x) &= \frac{x}{|x| + 2x^2} = \lim_{h \rightarrow 0} \frac{0 - h}{|0 - h| + 2(0 - h)^2} \\ &= \lim_{h \rightarrow 0} \frac{-h}{h + 2h^2} = \lim_{h \rightarrow 0} \frac{-h}{h(1 + 2h)} \\ &= \lim_{h \rightarrow 0} \frac{-1}{1 + 2h} = \frac{-1}{1 + 2(0)} = -1 \end{aligned}$$

$$\begin{aligned} \lim_{x \rightarrow 0^+} f(x) &= \frac{x}{|x| + 2x^2} = \lim_{h \rightarrow 0} \frac{0 + h}{|0 + h| + 2(0 + h)^2} \\ &= \lim_{h \rightarrow 0} \frac{h}{h + 2h^2} = \lim_{h \rightarrow 0} \frac{h}{h(1 + 2h)} = \frac{1}{1 + 0} = 1 \end{aligned}$$

$$\lim_{x \rightarrow 0^-} f(x) \neq \lim_{x \rightarrow 0^+} f(x)$$

Now, as the left hand limit and the right hand limit are not equal and the value of both the limits are a constant.

Hence, regardless the choice of  $k$ , the given function remains discontinuous at  $x = 0$ .

**16. Find the values of  $a$  and  $b$  such that the function  $f$  defined by**

$$f(x) = \begin{cases} \frac{x-4}{|x-4|} + a, & \text{if } x < 4 \\ a+b, & \text{if } x = 4 \\ \frac{x-4}{|x-4|} + b, & \text{if } x > 4 \end{cases}$$



is a continuous function at  $x = 4$ .

**Solution:**

Finding the left hand and right hand limit for the given function, we have

$$\begin{aligned}\lim_{x \rightarrow 4^-} f(x) &= \frac{x-4}{|x-4|} + a \\ &= \lim_{h \rightarrow 0} \frac{4-h-4}{|4-h-4|} + a = \lim_{h \rightarrow 0} \frac{-h}{h} + a = -1 + a \\ \lim_{x \rightarrow 4} f(x) &= a + b \\ \lim_{x \rightarrow 4^+} f(x) &= \frac{x-4}{|x-4|} + b \\ &= \lim_{h \rightarrow 0} \frac{4+h-4}{|4+h-4|} + b = \lim_{h \rightarrow 0} \frac{h}{h} + b = 1 + b\end{aligned}$$

As the function is continuous at  $x = 4$ .

$$\therefore \lim_{x \rightarrow 4^-} f(x) = \lim_{x \rightarrow 4} f(x) = \lim_{x \rightarrow 4^+} f(x)$$

$$\text{So, } -1 + a = a + b = 1 + b$$

$$-1 + a = a + b \text{ and } 1 + b = a + b$$

$$\text{We get, } b = -1 \text{ and } 1 + -1 = a + -1 \Rightarrow a = 1$$

Therefore, the value of  $a = 1$  and  $b = -1$

**17. Given the function  $f(x) = 1/(x+2)$ . Find the points of discontinuity of the composite function  $y = f(f(x))$ .**

**Solution:**

Given,

$$\begin{aligned}f(x) &= \frac{1}{x+2} \\ f[f(x)] &= \frac{1}{f(x)+2} = \frac{1}{\frac{1}{x+2}+2} = \frac{1}{\frac{1+2x+4}{x+2}} = \frac{x+2}{2x+5} \\ \therefore f[f(x)] &= \frac{x+2}{2x+5}\end{aligned}$$

Now, the function will not be defined and continuous where

$$2x + 5 = 0 \Rightarrow x = -5/2$$

Therefore,  $x = -5/2$  is the point of discontinuity.

**18. Find all points of discontinuity of the function  $f(t) = \frac{1}{t^2+t-2}$ , where  $t = \frac{1}{x-1}$ .**

**Solution:**

$$\begin{aligned}
 \text{Given, } f(t) &= \frac{1}{t^2 + t - 2} \\
 \Rightarrow f(t) &= \frac{1}{\frac{1}{(x-1)^2} + \frac{1}{(x-1)} - 2} \quad \left[ \text{putting } t = \frac{1}{x-1} \right] \\
 &= \frac{1}{\frac{1 + x - 1 - 2(x-1)^2}{(x-1)^2}} = \frac{(x-1)^2}{x - 2x^2 - 2 + 4x} \\
 &= \frac{(x-1)^2}{-2x^2 + 5x - 2} = \frac{(x-1)^2}{-(2x^2 - 5x + 2)} \\
 &= \frac{(x-1)^2}{-[2x^2 - 4x - x + 2]} = \frac{(x-1)^2}{-[2x(x-2) - 1(x-2)]} \\
 &= \frac{(x-1)^2}{-(x-2)(2x-1)} = \frac{(x-1)^2}{(2-x)(2x-1)}
 \end{aligned}$$

Now,

if  $f(t)$  is discontinuous, then  $2 - x = 0 \Rightarrow x = 2$

And,  $2x - 1 = 0 \Rightarrow x = \frac{1}{2}$

Therefore, the required points of discontinuity for the given function are 2 and  $\frac{1}{2}$ .

**19. Show that the function  $f(x) = |\sin x + \cos x|$  is continuous at  $x = p$ . Examine the differentiability of  $f$ , where  $f$  is defined by**

**Solution:**

Given,

$f(x) = |\sin x + \cos x|$  at  $x = \pi$

Now, put  $g(x) = \sin x + \cos x$  and  $h(x) = |x|$

Hence,  $h[g(x)] = h(\sin x + \cos x) = |\sin x + \cos x|$

Now,

$g(x) = \sin x + \cos x$  is a continuous function since  $\sin x$  and  $\cos x$  are two continuous functions at  $x = \pi$ .

We know that, every modulus function is a continuous function everywhere.

Therefore,  $f(x) = |\sin x + \cos x|$  is continuous function at  $x = \pi$ .

$$f(x) = \begin{cases} x[x], & \text{if } 0 \leq x < 2 \\ (x-1)x, & \text{if } 2 \leq x < 3 \end{cases}$$

**20. at  $x = 2$ .**

**Solution:**

We know that, a function  $f$  is differentiable at a point 'a' in its domain if

$Lf'(c) = Rf'(c)$

where  $Lf'(c) = \lim_{h \rightarrow 0} \frac{f(a-h) - f(a)}{-h}$  and

$$Rf'(c) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

Here,  $f(x) = \begin{cases} x[x], & \text{if } 0 \leq x < 2 \\ (x-1)x, & \text{if } 2 \leq x < 2 \end{cases}$  at  $x = 2$ .

$$\begin{aligned} Lf'(c) &= \lim_{h \rightarrow 0} \frac{f(2-h) - f(2)}{-h} = \lim_{h \rightarrow 0} \frac{(2-h)[2-h] - (2-1)2}{-h} \\ &= \lim_{h \rightarrow 0} \frac{(2-h) \cdot 1 - 2}{-h} \quad [\because [2-h] = 1] \\ &= \lim_{h \rightarrow 0} \frac{2-h-2}{-h} = 1 \end{aligned}$$

$$\begin{aligned} Rf'(c) &= \lim_{h \rightarrow 0} \frac{f(2+h) - f(2)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(2+h-1)(2+h) - (2-1) \cdot 2}{h} \\ &= \lim_{h \rightarrow 0} \frac{(1+h)(2+h) - 2}{h} = \lim_{h \rightarrow 0} \frac{2+h+2h+h^2-2}{h} \\ &= \lim_{h \rightarrow 0} \frac{3h+h^2}{h} = \lim_{h \rightarrow 0} \frac{h(3+h)}{h} = 3 \end{aligned}$$

$$Lf'(2) \neq Rf'(2)$$

Therefore,  $f(x)$  is not differentiable at  $x = 2$ .

$$21. \quad f(x) = \begin{cases} x^2 \sin \frac{1}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

at  $x = 0$ .

**Solution:**

Given,

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases} \quad \text{at } x = 0$$

For differentiability we know that:

$$Lf'(c) = Rf'(c)$$

$$\therefore Lf'(0) = \lim_{h \rightarrow 0} \frac{f(0-h) - f(0)}{-h}$$

$$\begin{aligned}
 &= \lim_{h \rightarrow 0} \frac{(0-h)^2 \sin \frac{1}{(0-h)} - 0}{-h} = \frac{h^2 \cdot \sin\left(-\frac{1}{h}\right)}{-h} \\
 &= h \cdot \sin\left(\frac{1}{h}\right) = 0 \times \left[-1 \leq \sin\left(\frac{1}{h}\right) \leq 1\right] \\
 &= 0
 \end{aligned}$$

$$\begin{aligned}
 Rf'(0) &= \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{(0+h)^2 \sin\left(\frac{1}{0+h}\right) - 0}{h} \\
 &= \lim_{h \rightarrow 0} \frac{h^2 \sin\left(\frac{1}{h}\right)}{h} = \lim_{h \rightarrow 0} h \cdot \sin\left(\frac{1}{h}\right) \\
 &= 0 \times \left[-1 \leq \sin\left(\frac{1}{h}\right) \leq 1\right] = 0
 \end{aligned}$$

Hence,  $Lf'(0) = Rf'(0) = 0$

Therefore,  $f(x)$  is differentiable at  $x = 0$

22.  $f(x) = \begin{cases} 1+x & , \text{ if } x \leq 2 \\ 5-x & , \text{ if } x > 2 \end{cases}$   
at  $x = 2$ .

**Solution:**

We know that,  $f(x)$  is differentiable at  $x = 2$  if

$$Lf'(2) = Rf'(2)$$

Now,

$$\begin{aligned}
 Lf'(2) &= \lim_{h \rightarrow 0} \frac{f(2-h) - f(2)}{-h} \\
 &= \lim_{h \rightarrow 0} \frac{(1+2-h) - (1+2)}{-h} = \lim_{h \rightarrow 0} \frac{3-h-3}{-h} = \frac{-h}{-h} = 1
 \end{aligned}$$

$$\begin{aligned}
 Rf'(2) &= \lim_{h \rightarrow 0} \frac{f(2+h) - f(2)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{[5-(2+h)] - (1+2)}{h} = \lim_{h \rightarrow 0} \frac{3-h-3}{h} \\
 &= \frac{-h}{h} = -1
 \end{aligned}$$

$$\text{So, } Lf'(2) \neq Rf'(2)$$

Thus,  $f(x)$  is not differentiable at  $x = 2$ .

**23. Show that  $f(x) = |x - 5|$  is continuous but not differentiable at  $x = 5$ .**

**Solution:**



Given,  $f(x) = |x - 5|$

$$\Rightarrow f(x) = \begin{cases} -(x - 5) & \text{if } x - 5 < 0 \text{ or } x < 5 \\ x - 5 & \text{if } x - 5 > 0 \text{ or } x > 5 \end{cases}$$

For continuity at  $x = 5$

$$\begin{aligned} \text{L.H.L. } \lim_{x \rightarrow 5^-} f(x) &= -(x - 5) \\ &= \lim_{h \rightarrow 0} -(5 - h - 5) = \lim_{h \rightarrow 0} h = 0 \end{aligned}$$

$$\begin{aligned} \text{R.H.L. } \lim_{x \rightarrow 5^+} f(x) &= x - 5 \\ &= \lim_{h \rightarrow 0} (5 + h - 5) = \lim_{h \rightarrow 0} h = 0 \end{aligned}$$

L.H.L. = R.H.L.

So,  $f(x)$  is continuous at  $x = 5$ .

Now, for differentiability

$$\begin{aligned} \text{Lf}'(5) &= \lim_{h \rightarrow 0} \frac{f(5 - h) - f(5)}{-h} \\ &= \lim_{h \rightarrow 0} \frac{-(5 - h - 5) - (5 - 5)}{-h} = \lim_{h \rightarrow 0} \frac{h}{-h} = -1 \end{aligned}$$

$$\begin{aligned} \text{Rf}'(5) &= \lim_{h \rightarrow 0} \frac{f(5 + h) - f(5)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(5 + h - 5) - (5 - 5)}{h} = \lim_{h \rightarrow 0} \frac{h - 0}{h} = 1 \end{aligned}$$

Thus,  $\text{Lf}'(5) \neq \text{Rf}'(5)$

Therefore,  $f(x)$  is not differentiable at  $x = 5$ .

**24. A function  $f: \mathbb{R} \rightarrow \mathbb{R}$  satisfies the equation  $f(x + y) = f(x)f(y)$  for all  $x, y \in \mathbb{R}$ ,  $f(x) \neq 0$ .**

**Suppose that the function is differentiable at  $x = 0$  and  $f'(0) = 2$ . Prove that  $f'(x) = 2f(x)$ .**

**Solution:**

Given,

$f: \mathbb{R} \rightarrow \mathbb{R}$  satisfies the equation  $f(x + y) = f(x)f(y)$  for all  $x, y \in \mathbb{R}$ ,  $f(x) \neq 0$

Let us take any point  $x = 0$  at which the function  $f(x)$  is differentiable.

$$\begin{aligned} \text{So, } f'(0) &= \lim_{h \rightarrow 0} \frac{f(0 + h) - f(0)}{h} \\ 2 &= \lim_{h \rightarrow 0} \frac{f(0) \cdot f(h) - f(0)}{h} \quad [\because f(0) = f(h)] \quad \dots(i) \end{aligned}$$

$$\Rightarrow 2 = \lim_{h \rightarrow 0} \frac{f(0)[f(h) - 1]}{h}$$

$$\begin{aligned} \text{Now, } f'(x) &= \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x) \cdot f(h) - f(x)}{h} \quad [\because f(x + y) = f(x) \cdot f(y)] \end{aligned}$$

$$= \lim_{h \rightarrow 0} \frac{f(x)[f(h)-1]}{h} = 2f(x) \quad \text{from eqn. (i)}$$

Therefore,  $f'(x) = 2f(x)$ .

**Differentiate each of the following w.r.t.  $x$  (Exercises 25 to 43) :**

25.  $2^{\cos^2 x}$

**Solution:**

Let  $y = 2^{\cos^2 x}$

Taking log on both sides, we get

$$\log y = \log 2^{\cos^2 x} \Rightarrow \log y = \cos^2 x \cdot \log 2$$

Now,

Differentiating both sides w.r.t.  $x$

$$\frac{1}{y} \cdot \frac{dy}{dx} = \log 2 \cdot \frac{d}{dx} \cos^2 x$$

$$\frac{1}{y} \cdot \frac{dy}{dx} = \log 2 \left[ 2 \cos x \cdot \frac{d}{dx} \cos x \right]$$

$$\frac{1}{y} \cdot \frac{dy}{dx} = \log 2 [2 \cos x (-\sin x)]$$

$$\frac{1}{y} \cdot \frac{dy}{dx} = \log 2 (-\sin 2x)$$

$$\frac{dy}{dx} = -y \cdot \log 2 \sin 2x$$

Thus,  $\frac{dy}{dx} = -2^{\cos^2 x} (\log 2 \sin 2x)$

26.  $\frac{8^x}{x^8}$

**Solution:**

Let  $y = \frac{8^x}{x^8}$

Taking log on both sides, we get  $\log y = \log \frac{8^x}{x^8}$

$$\Rightarrow \log y = \log 8^x - \log x^8 \Rightarrow \log y = x \log 8 - 8 \log x$$

Differentiating both sides w.r.t.  $x$

$$\frac{1}{y} \cdot \frac{dy}{dx} = \log 8 - \frac{8}{x} \Rightarrow \frac{dy}{dx} = y \left[ \log 8 - \frac{8}{x} \right]$$

Thus,  $\frac{dy}{dx} = \frac{8^x}{x^8} \left[ \log 8 - \frac{8}{x} \right]$

**NAGEEN GROUP**  
tamso ma jyotirgamaya

27.  $\log \left( x + \sqrt{x^2 + a} \right)$

**Solution:**

Let  $y = \log \left( x + \sqrt{x^2 + a} \right)$

Differentiating both sides w.r.t.  $x$

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} \log \left( x + \sqrt{x^2 + a} \right) \\ &= \frac{1}{x + \sqrt{x^2 + a}} \cdot \frac{d}{dx} \left( x + \sqrt{x^2 + a} \right) \\ &= \frac{1}{x + \sqrt{x^2 + a}} \cdot \left[ 1 + \frac{1}{2\sqrt{x^2 + a}} \times \frac{d}{dx} (x^2 + a) \right] \\ &= \frac{1}{x + \sqrt{x^2 + a}} \cdot \left[ 1 + \frac{1}{2\sqrt{x^2 + a}} \cdot 2x \right] \\ &= \frac{1}{x + \sqrt{x^2 + a}} \cdot \left[ 1 + \frac{x}{\sqrt{x^2 + a}} \right] \\ &= \frac{1}{x + \sqrt{x^2 + a}} \cdot \left( \frac{\sqrt{x^2 + a} + x}{\sqrt{x^2 + a}} \right) = \frac{1}{\sqrt{x^2 + a}} \end{aligned}$$

Thus,  $\frac{dy}{dx} = \frac{1}{\sqrt{x^2 + a}}$

28.  $\log \left[ \log (\log x^5) \right]$

**Solution:**

Let,  $y = \log [\log (\log x^5)]$

Differentiating both sides w.r.t.  $x$

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} \log [\log (\log x^5)] \\ &= \frac{1}{\log (\log x^5)} \times \frac{d}{dx} \log (\log x^5) \\ &= \frac{1}{\log (\log x^5)} \times \frac{1}{\log (x^5)} \times \frac{d}{dx} \log x^5 \\ &= \frac{1}{\log (\log x^5)} \cdot \frac{1}{\log (x^5)} \cdot \frac{1}{x^5} \cdot \frac{d}{dx} x^5 \end{aligned}$$

$$= \frac{1}{\log(\log x^5)} \cdot \frac{1}{\log(x^5)} \cdot \frac{1}{x^5} \cdot 5x^4$$

$$= \frac{5}{x \log(x^5) \cdot \log(\log x^5)}$$

Thus,  $\frac{dy}{dx} = \frac{5}{x \log(x^5) \cdot \log(\log x^5)}$

29.  $\sin \sqrt{x} + \cos^2 \sqrt{x}$

**Solution:**

Let  $y = \sin \sqrt{x} + \cos^2 \sqrt{x}$

Differentiating both sides w.r.t.  $x$

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx}(\sin \sqrt{x}) + \frac{d}{dx}(\cos^2 \sqrt{x}) \\ &= \cos \sqrt{x} \cdot \frac{d}{dx}(\sqrt{x}) + 2 \cos \sqrt{x} \cdot \frac{d}{dx}(\cos \sqrt{x}) \\ &= \cos \sqrt{x} \cdot \frac{1}{2\sqrt{x}} + 2 \cos \sqrt{x} (-\sin \sqrt{x}) \cdot \frac{d}{dx} \sqrt{x} \\ &= \frac{1}{2\sqrt{x}} \cdot \cos \sqrt{x} - 2 \cos \sqrt{x} \cdot \sin \sqrt{x} \cdot \frac{1}{2\sqrt{x}} \\ &= \frac{\cos \sqrt{x}}{2\sqrt{x}} - \frac{\sin 2\sqrt{x}}{2\sqrt{x}} \end{aligned}$$

Thus,  $\frac{dy}{dx} = \frac{\cos \sqrt{x}}{2\sqrt{x}} - \frac{\sin 2\sqrt{x}}{2\sqrt{x}}$

30.  $\sin^n(ax^2 + bx + c)$

**Solution:**

Let  $y = \sin^n(ax^2 + bx + c)$

Differentiating both sides w.r.t.  $x$

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} \sin^n(ax^2 + bx + c) \\ &= n \cdot \sin^{n-1}(ax^2 + bx + c) \cdot \frac{d}{dx} \sin(ax^2 + bx + c) \\ &= n \cdot \sin^{n-1}(ax^2 + bx + c) \cdot \cos(ax^2 + bx + c) \cdot \frac{d}{dx} (ax^2 + bx + c) \\ &= n \cdot \sin^{n-1}(ax^2 + bx + c) \cdot \cos(ax^2 + bx + c) \cdot (2ax + b) \end{aligned}$$

Thus,  $\frac{dy}{dx} = n(2ax + b) \cdot \sin^{n-1}(ax^2 + bx + c) \cdot \cos(ax^2 + bx + c)$



31.  $\cos(\tan \sqrt{x+1})$

**Solution:**

Let  $y = \cos(\tan \sqrt{x+1})$

Differentiating both sides w.r.t.  $x$

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} \cos(\tan \sqrt{x+1}) \\ &= -\sin(\tan \sqrt{x+1}) \cdot \frac{d}{dx}(\tan \sqrt{x+1}) \\ &= -\sin(\tan \sqrt{x+1}) \cdot \sec^2 \sqrt{x+1} \cdot \frac{d}{dx} \sqrt{x+1} \\ &= -\sin(\tan \sqrt{x+1}) \cdot \sec^2 \sqrt{x+1} \cdot \frac{1}{2\sqrt{x+1}} \cdot 1\end{aligned}$$

Thus,  $\frac{dy}{dx} = -\frac{1}{2\sqrt{x+1}} \sin(\tan \sqrt{x+1}) \cdot \sec^2 \sqrt{x+1}$

32.  $\sin x^2 + \sin^2 x + \sin^2(x^2)$

**Solution:**

Let  $y = \sin x^2 + \sin^2 x + \sin^2(x^2)$

Differentiating both sides w.r.t.  $x$ ,

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx} \sin(x^2) + \frac{d}{dx} \sin^2 x + \frac{d}{dx} \sin^2(x^2) \\ &= \cos x^2 \cdot \frac{d}{dx}(x^2) + 2 \sin x \cdot \frac{d}{dx}(\sin x) + 2 \sin(x^2) \cdot \frac{d}{dx} \sin(x^2) \\ &= \cos x^2 \cdot 2x + 2 \sin x \cdot \cos x + 2 \sin x^2 \cdot \cos x^2 \cdot \frac{d}{dx}(x^2) \\ &= 2x \cdot \cos x^2 + \sin 2x + 2 \sin x^2 \cdot \cos x^2 \cdot 2x\end{aligned}$$

Thus,  $\frac{dy}{dx} = 2x \cdot \cos x^2 + \sin 2x + 2x \sin 2x^2$

33.  $\sin^{-1}\left(\frac{1}{\sqrt{x+1}}\right)$

**Solution:**

Let  $y = \sin^{-1}\left(\frac{1}{\sqrt{x+1}}\right)$

Differentiating both sides w.r.t.  $x$

$$\frac{dy}{dx} = \frac{d}{dx} \sin^{-1}\left(\frac{1}{\sqrt{x+1}}\right) = \frac{1}{\sqrt{1-\left(\frac{1}{\sqrt{x+1}}\right)^2}} \cdot \frac{d}{dx}\left(\frac{1}{\sqrt{x+1}}\right)$$

$$= \frac{1}{\sqrt{1-\frac{1}{x+1}}} \cdot \frac{d}{dx}(x+1)^{-1/2}$$

$$= \frac{1}{\sqrt{\frac{x+1-1}{x+1}}} \cdot \frac{-1}{2}(x+1)^{-3/2} \cdot \frac{d}{dx}(x+1)$$

$$= \frac{\sqrt{x+1}}{\sqrt{x}} \cdot \frac{-1}{2}(x+1)^{-3/2} \cdot 1$$

$$= \frac{-1}{2} \cdot \frac{\sqrt{x+1}}{\sqrt{x}} \cdot \frac{1}{(x+1)^{3/2}} = -\frac{1}{2\sqrt{x}(x+1)}$$

Thus,  $\frac{dy}{dx} = -\frac{1}{2\sqrt{x}(x+1)}$

### 34. $(\sin x)^{\cos x}$

**Solution:**

Let  $y = (\sin x)^{\cos x}$

Taking log on both sides,

$$\log y = \log (\sin x)^{\cos x}$$

$$\Rightarrow \log y = \cos x \cdot \log (\sin x)$$

$$[\because \log x^y = y \log x]$$

Differentiating both sides w.r.t.  $x$ ,

$$\frac{1}{y} \cdot \frac{dy}{dx} = \frac{d}{dx} \cos x \cdot \log (\sin x)$$

$$\frac{1}{y} \cdot \frac{dy}{dx} = \cos x \cdot \frac{d}{dx} \log (\sin x) + \log (\sin x) \cdot \frac{d}{dx} \cos x$$

$$\frac{1}{y} \cdot \frac{dy}{dx} = \cos x \cdot \frac{1}{\sin x} \cdot \frac{d}{dx} (\sin x) + \log (\sin x) \cdot (-\sin x)$$

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = \cot x \cdot \cos x - \sin x \cdot \log (\sin x)$$

$$\frac{dy}{dx} = y [\cot x \cdot \cos x - \sin x \cdot \log (\sin x)]$$

$$\text{Thus, } \frac{dy}{dx} = (\sin x)^{\cos x} \left[ \frac{\cos^2 x}{\sin x} - \sin x \cdot \log(\sin x) \right]$$

### 35. $\sin^m x \cdot \cos^n x$

**Solution:**

$$\text{Let } y = \sin^m x \cdot \cos^n x$$

Differentiating both sides w.r.t.  $x$

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx}(\sin^m x \cdot \cos^n x) \\ &= \sin^m x \cdot \frac{d}{dx}(\cos^n x) + \cos^n x \cdot \frac{d}{dx} \sin^m x \\ &= \sin^m x \cdot n \cdot \cos^{n-1} x \frac{d}{dx}(\cos x) + \cos^n x \cdot m \cdot \sin^{m-1} x \frac{d}{dx}(\sin x) \\ &= n \cdot \sin^m x \cdot \cos^{n-1} x \cdot (-\sin x) + m \cdot \cos^n x \cdot \sin^{m-1} x \cdot \cos x \\ &= -n \cdot \sin^{m+1} x \cdot \cos^{n-1} x + m \cdot \cos^{n+1} x \cdot \sin^{m-1} x \\ &= \sin^m x \cdot \cos^n x \left[ -n \frac{\sin x}{\cos x} + m \frac{\cos x}{\sin x} \right] \end{aligned}$$

$$\text{Thus, } \frac{dy}{dx} = \sin^m x \cdot \cos^n x [-n \tan x + m \cot x]$$

### 36. $(x+1)^2 + (x+2)^3 + (x+3)^4$

**Solution:**

$$\text{Let } y = (x+1)^2(x+2)^3(x+3)^4$$

Taking log on both sides,

$$\log y = \log [(x+1)^2 \cdot (x+2)^3 \cdot (x+3)^4]$$

$$\log y = \log (x+1)^2 + \log (x+2)^3 + \log (x+3)^4$$

$$[\because \log xy = \log x + \log y]$$

$$\Rightarrow \log y = 2 \log (x+1) + 3 \log (x+2) + 4 \log (x+3)$$

$$[\because \log x^y = y \log x]$$

Differentiating both sides w.r.t.  $x$ ,

$$\frac{1}{y} \cdot \frac{dy}{dx} = 2 \cdot \frac{d}{dx} \log(x+1) + 3 \cdot \frac{d}{dx} \log(x+2) + 4 \cdot \frac{d}{dx} \log(x+3)$$

$$\frac{1}{y} \cdot \frac{dy}{dx} = 2 \cdot \frac{1}{x+1} + 3 \cdot \frac{1}{x+2} + 4 \cdot \frac{1}{x+3}$$

$$\frac{dy}{dx} = y \left[ \frac{2}{x+1} + \frac{3}{x+2} + \frac{4}{x+3} \right]$$

Now,

$$\begin{aligned}\Rightarrow \frac{dy}{dx} &= (x+1)^2(x+2)^3(x+3)^4 \left[ \frac{2}{x+1} + \frac{3}{x+2} + \frac{4}{x+3} \right] \\ &= (x+1)^2(x+2)^3(x+3)^4 \\ &\quad \left[ \frac{2(x+2)(x+3) + 3(x+1)(x+3) + 4(x+1)(x+2)}{(x+1)(x+2)(x+3)} \right] \\ &= (x+1)(x+2)^2(x+3)^3(2x^2 + 10x + 12 + 3x^2 + 12x + 9 \\ &\quad + 4x^2 + 12x + 8) \\ &= (x+1)(x+2)^2(x+3)^3(9x^2 + 34x + 29)\end{aligned}$$

$$\text{Thus, } \frac{dy}{dx} = (x+1)(x+2)^2(x+3)^3(9x^2 + 34x + 29)$$

$$\cos^{-1}\left(\frac{\sin x + \cos x}{\sqrt{2}}\right), \frac{-\pi}{4} < x < \frac{\pi}{4}$$

37.

**Solution:**

$$\begin{aligned}\text{Let } y &= \cos^{-1}\left(\frac{\sin x + \cos x}{\sqrt{2}}\right) \\ &= \cos^{-1}\left[\frac{1}{\sqrt{2}} \sin x + \frac{1}{\sqrt{2}} \cos x\right] \\ &= \cos^{-1}\left[\sin \frac{\pi}{4} \sin x + \cos \frac{\pi}{4} \cdot \cos x\right] = \cos^{-1}\left[\cos\left(\frac{\pi}{4} - x\right)\right] \\ y &= \frac{\pi}{4} - x \quad \left[\because -\frac{\pi}{4} < x < \frac{\pi}{4}\right]\end{aligned}$$

Differentiating both sides w.r.t.  $x$

$$\text{Thus, } \frac{dy}{dx} = -1$$

$$\tan^{-1}\left(\frac{\sqrt{1-\cos x}}{\sqrt{1+\cos x}}\right), -\frac{\pi}{4} < x < \frac{\pi}{4}$$

38.

**Solution:**

$$\begin{aligned}\text{Let } y &= \tan^{-1}\left[\sqrt{\frac{1-\cos x}{1+\cos x}}\right] \\ &= \tan^{-1}\left[\sqrt{\frac{2\sin^2 x/2}{2\cos^2 x/2}}\right] \quad \left[\because 1-\cos x = 2\sin^2 x/2\right. \\ &\quad \left.1+\cos x = 2\cos^2 x/2\right]\end{aligned}$$



$$= \tan^{-1} \left[ \frac{\sin x/2}{\cos x/2} \right] = \tan^{-1} \left[ \tan \frac{x}{2} \right]$$

Thus,  $y = \frac{x}{2}$

Differentiating both sides w.r.t.  $x$

$$\frac{dy}{dx} = \frac{1}{2} \frac{d}{dx}(x) = \frac{1}{2} \cdot 1 = \frac{1}{2}$$

Hence,  $\frac{dy}{dx} = \frac{1}{2}$

$$\tan^{-1}(\sec x + \tan x), -\frac{\pi}{2} < x < \frac{\pi}{2}$$

39.

**Solution:**

Let  $y = \tan^{-1}(\sec x + \tan x)$

Differentiating both sides w.r.t.  $x$

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} [\tan^{-1}(\sec x + \tan x)] \\ &= \frac{1}{1 + (\sec x + \tan x)^2} \cdot \frac{d}{dx} (\sec x + \tan x) \\ &= \frac{1}{1 + \sec^2 x + \tan^2 x + 2 \sec x \tan x} \cdot (\sec x \tan x + \sec^2 x) \\ &= \frac{1}{(1 + \tan^2 x) + \sec^2 x + 2 \sec x \tan x} \cdot \sec x (\tan x + \sec x) \\ &= \frac{1}{\sec^2 x + \sec^2 x + 2 \sec x \tan x} \cdot \sec x (\tan x + \sec x) \\ &= \frac{1}{2 \sec^2 x + 2 \sec x \tan x} \cdot \sec x (\tan x + \sec x) \\ &= \frac{1}{2 \sec x (\sec x + \tan x)} \cdot \sec x (\tan x + \sec x) = \frac{1}{2} \end{aligned}$$

Thus,  $\frac{dy}{dx} = \frac{1}{2}$

$$\tan^{-1} \left( \frac{a \cos x - b \sin x}{b \cos x + a \sin x} \right), -\frac{\pi}{2} < x < \frac{\pi}{2} \text{ and } \frac{a}{b} \tan x > -1$$

40.

**Solution:**



$$\text{Let } y = \tan^{-1} \left( \frac{a \cos x - b \sin x}{b \cos x + a \sin x} \right)$$

$$y = \tan^{-1} \left[ \frac{\frac{a \cos x}{b \cos x} - \frac{b \sin x}{b \cos x}}{\frac{b \cos x}{b \cos x} + \frac{a \sin x}{b \cos x}} \right]$$

$$y = \tan^{-1} \left[ \frac{\frac{a}{b} - \tan x}{1 + \frac{a}{b} \tan x} \right]$$

$$y = \tan^{-1} \frac{a}{b} - \tan^{-1}(\tan x)$$

$$\left[ \because \tan^{-1} \left( \frac{x-y}{1+xy} \right) = \tan^{-1} x - \tan^{-1} y \right]$$

$$\Rightarrow y = \tan^{-1} \frac{a}{b} - x$$

Differentiating both sides with respect to  $x$

$$\frac{dy}{dx} = \frac{d}{dx} \left( \tan^{-1} \frac{a}{b} \right) - \frac{d}{dx}(x) = 0 - 1 = -1$$

Thus,  $\frac{dy}{dx} = -1$ .

$$\sec^{-1} \left( \frac{1}{4x^3 - 3x} \right), 0 < x < \frac{1}{\sqrt{2}}$$

41.

**Solution:**

$$\text{Let } y = \sec^{-1} \left( \frac{1}{4x^3 - 3x} \right)$$

Put  $x = \cos \theta \therefore \theta = \cos^{-1} x$

$$\Rightarrow y = \sec^{-1} \left( \frac{1}{4 \cos^3 \theta - 3 \cos \theta} \right)$$

$$y = \sec^{-1} \left( \frac{1}{\cos 3\theta} \right) \quad [\because \cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta]$$

$$y = \sec^{-1}(\sec 3\theta) \Rightarrow y = 3\theta$$

$$y = 3 \cos^{-1} x$$

Differentiating both sides w.r.t.  $x$

$$\frac{dy}{dx} = 3 \cdot \frac{d}{dx} \cos^{-1} x = 3 \left( \frac{-1}{\sqrt{1-x^2}} \right) = \frac{-3}{\sqrt{1-x^2}}$$

Thus,  $\frac{dy}{dx} = \frac{-3}{\sqrt{1-x^2}}$ .

$$\tan^{-1} \left( \frac{3a^2x - x^3}{a^3 - 3ax^2} \right), \frac{-1}{\sqrt{3}} < \frac{x}{a} < \frac{1}{\sqrt{3}}$$

42.

**Solution:**

$$\text{Let } y = \tan^{-1} \left[ \frac{3a^2x - x^3}{a^3 - 3ax^2} \right]$$

$$\text{Put } x = a \tan \theta \quad \therefore \theta = \tan^{-1} \frac{x}{a}$$

$$y = \tan^{-1} \left[ \frac{3a^2 \cdot a \tan \theta - a^3 \tan^3 \theta}{a^3 - 3a \cdot a^2 \tan^2 \theta} \right]$$

$$y = \tan^{-1} \left[ \frac{3a^3 \tan \theta - a^3 \tan^3 \theta}{a^3 - 3a^3 \tan^2 \theta} \right]$$

$$y = \tan^{-1} \left[ \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta} \right]$$

$$y = \tan^{-1} [\tan 3\theta] \quad \left[ \because \tan 3\theta = \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta} \right]$$

$$\Rightarrow y = 3\theta \Rightarrow y = 3 \tan^{-1} \frac{x}{a}$$

Differentiating both sides w.r.t.  $x$

$$\begin{aligned} \frac{dy}{dx} &= 3 \cdot \frac{d}{dx} \left( \tan^{-1} \frac{x}{a} \right) \\ &= 3 \cdot \frac{1}{1 + \frac{x^2}{a^2}} \cdot \frac{d}{dx} \left( \frac{x}{a} \right) = 3 \cdot \frac{a^2}{a^2 + x^2} \cdot \frac{1}{a} = \frac{3a}{a^2 + x^2} \end{aligned}$$

$$\text{Thus, } \frac{dy}{dx} = \frac{3a}{a^2 + x^2}.$$

$$\tan^{-1} \left( \frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}} \right), -1 < x < 1, x \neq 0$$

43.

**Solution:**

$$\text{Let } y = \tan^{-1} \left( \frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}} \right)$$

$$\text{Putting } x^2 = \cos 2\theta \quad \therefore \theta = \frac{1}{2} \cos^{-1} x^2$$

$$y = \tan^{-1} \left( \frac{\sqrt{1+\cos 2\theta} + \sqrt{1-\cos 2\theta}}{\sqrt{1+\cos 2\theta} - \sqrt{1-\cos 2\theta}} \right)$$

$$y = \tan^{-1} \left( \frac{\sqrt{2 \cos^2 \theta} + \sqrt{2 \sin^2 \theta}}{\sqrt{2 \cos^2 \theta} - \sqrt{2 \sin^2 \theta}} \right)$$

$$y = \tan^{-1} \left( \frac{\sqrt{2} \cos \theta + \sqrt{2} \sin \theta}{\sqrt{2} \cos \theta - \sqrt{2} \sin \theta} \right)$$

$$y = \tan^{-1} \left( \frac{\cos \theta + \sin \theta}{\cos \theta - \sin \theta} \right)$$

$$y = \tan^{-1} \left[ \frac{\cos \theta + \sin \theta}{\cos \theta - \sin \theta} \right]$$

$$y = \tan^{-1} \left[ \frac{1 + \tan \theta}{1 - \tan \theta} \right]$$

$$y = \tan^{-1} \left[ \frac{\tan \frac{\pi}{4} + \tan \theta}{1 - \tan \frac{\pi}{4} \cdot \tan \theta} \right]$$

$$y = \tan^{-1} \left[ \tan \left( \frac{\pi}{4} + \theta \right) \right]$$

$$\Rightarrow y = \frac{\pi}{4} + \theta \Rightarrow y = \frac{\pi}{4} + \frac{1}{2} \cos^{-1} x^2$$

Differentiating both sides w.r.t.  $x$

$$\frac{dy}{dx} = \frac{d}{dx} \left( \frac{\pi}{4} \right) + \frac{1}{2} \frac{d}{dx} (\cos^{-1} x^2)$$

$$= 0 + \frac{1}{2} \times \frac{-1}{\sqrt{1-x^4}} \cdot \frac{d}{dx} (x^2) = \frac{-1.2x}{2\sqrt{1-x^4}} = -\frac{x}{\sqrt{1-x^4}}$$

$$\text{Thus, } \frac{dy}{dx} = -\frac{x}{\sqrt{1-x^4}}$$

Find  $dy/dx$  of each of the functions expressed in parametric form in Exercises from 44 to 48.

44.  $x = t + \frac{1}{t}, y = t - \frac{1}{t}$

**Solution:**

Given,

$$x = t + 1/t, y = t - 1/t$$

Differentiating both the parametric functions w.r.t  $\theta$

$$\frac{dx}{dt} = 1 - \frac{1}{t^2}, \frac{dy}{dt} = 1 + \frac{1}{t^2}$$

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{1 + \frac{1}{t^2}}{1 - \frac{1}{t^2}} = \frac{t^2 + 1}{t^2 - 1}$$

$$\text{Thus, } \frac{dy}{dx} = \frac{t^2 + 1}{t^2 - 1}$$

45.  $x = e^\theta \left( \theta + \frac{1}{\theta} \right), y = e^{-\theta} \left( \theta - \frac{1}{\theta} \right)$

**Solution:**

Given,

$$x = e^\theta \left( \theta + \frac{1}{\theta} \right), y = e^{-\theta} \left( \theta - \frac{1}{\theta} \right)$$

Differentiating both the parametric functions w.r.t.  $\theta$ .

$$\frac{dx}{d\theta} = e^\theta \left( 1 - \frac{1}{\theta^2} \right) + \left( \theta + \frac{1}{\theta} \right) \cdot e^\theta$$

$$\frac{dx}{d\theta} = e^\theta \left( 1 - \frac{1}{\theta^2} + \theta + \frac{1}{\theta} \right) \Rightarrow e^\theta \left( \frac{\theta^2 - 1 + \theta^3 + \theta}{\theta^2} \right)$$

$$= \frac{e^\theta (\theta^3 + \theta^2 + \theta - 1)}{\theta^2}$$

$$y = e^{-\theta} \left( \theta - \frac{1}{\theta} \right)$$

$$\frac{dy}{d\theta} = e^{-\theta} \left( 1 + \frac{1}{\theta^2} \right) + \left( \theta - \frac{1}{\theta} \right) \cdot (-e^{-\theta})$$

$$\frac{dy}{d\theta} = e^{-\theta} \left( 1 + \frac{1}{\theta^2} - \theta + \frac{1}{\theta} \right) \Rightarrow e^{-\theta} \left( \frac{\theta^2 + 1 - \theta^3 + \theta}{\theta^2} \right)$$

$$= e^{-\theta} \frac{(-\theta^3 + \theta^2 + \theta + 1)}{\theta^2}$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{dy/d\theta}{dx/d\theta} = \frac{e^{-\theta} \left( \frac{-\theta^3 + \theta^2 + \theta + 1}{\theta^2} \right)}{e^{\theta} \left( \frac{\theta^3 + \theta^2 + \theta - 1}{\theta^2} \right)} \\ &= e^{-2\theta} \left( \frac{-\theta^3 + \theta^2 + \theta + 1}{\theta^3 + \theta^2 + \theta - 1} \right) \\ \text{Thus, } \frac{dy}{dx} &= e^{-2\theta} \left( \frac{-\theta^3 + \theta^2 + \theta + 1}{\theta^3 + \theta^2 + \theta - 1} \right).\end{aligned}$$

**46.**  $x = 3\cos\theta - 2\cos^3\theta$ ,  $y = 3\sin\theta - 2\sin^3\theta$ .

**Solution:**

Given,  $x = 3\cos\theta - 2\cos^3\theta$ ,  $y = 3\sin\theta - 2\sin^3\theta$ .

Differentiating both the parametric functions w.r.t.  $\theta$

$$\begin{aligned}\frac{dx}{d\theta} &= -3\sin\theta - 6\cos^2\theta \cdot \frac{d}{d\theta}(\cos\theta) \\ &= -3\sin\theta - 6\cos^2\theta \cdot (-\sin\theta) \\ &= -3\sin\theta + 6\cos^2\theta \cdot \sin\theta \\ \frac{dy}{d\theta} &= 3\cos\theta - 6\sin^2\theta \cdot \frac{d}{d\theta}(\sin\theta) \\ &= 3\cos\theta - 6\sin^2\theta \cdot \cos\theta \\ \therefore \frac{dy}{dx} &= \frac{dy/d\theta}{dx/d\theta} = \frac{3\cos\theta - 6\sin^2\theta \cos\theta}{-3\sin\theta + 6\cos^2\theta \cdot \sin\theta} \\ \Rightarrow \frac{dy}{dx} &= \frac{\cos\theta(3 - 6\sin^2\theta)}{\sin\theta(-3 + 6\cos^2\theta)} = \frac{\cos\theta[3 - 6(1 - \cos^2\theta)]}{\sin\theta[-3 + 6\cos^2\theta]} \\ &= \cot\theta \left( \frac{3 - 6 + 6\cos^2\theta}{-3 + 6\cos^2\theta} \right) = \cot\theta \left( \frac{-3 + 6\cos^2\theta}{-3 + 6\cos^2\theta} \right) \\ &= \cot\theta\end{aligned}$$

Thus,  $\frac{dy}{dx} = \cot\theta$ .

**47.**  $\sin x = \frac{2t}{1+t^2}$ ,  $\tan y = \frac{2t}{1-t^2}$ .

**Solution:**

Given,

$\sin x = 2t/(1+t^2)$ ,  $\tan y = 2t/(1-t^2)$



Now, taking  $\sin x = \frac{2t}{1+t^2}$

Differentiating both sides w.r.t  $t$ , we get

$$\cos x \cdot \frac{dx}{dt} = \frac{(1+t^2) \cdot \frac{d}{dt}(2t) - 2t \cdot \frac{d}{dt}(1+t^2)}{(1+t^2)^2}$$

$$\cos x \cdot \frac{dx}{dt} = \frac{2(1+t^2) - 2t \cdot 2t}{(1+t^2)^2}$$

$$\frac{dx}{dt} = \frac{2+2t^2-4t^2}{(1+t^2)^2} \times \frac{1}{\cos x}$$

$$\frac{dx}{dt} = \frac{2-2t^2}{(1+t^2)^2} \times \frac{1}{\sqrt{1-\sin^2 x}}$$

$$\frac{dx}{dt} = \frac{2(1-t^2)}{(1+t^2)^2} \times \frac{1}{\sqrt{1-\left(\frac{2t}{1+t^2}\right)^2}}$$

$$\frac{dx}{dt} = \frac{2(1-t^2)}{(1+t^2)^2} \times \frac{1}{\sqrt{\frac{(1+t^2)^2-4t^2}{(1+t^2)^2}}}$$

$$\frac{dx}{dt} = \frac{2(1-t^2)}{(1+t^2)^2} \times \frac{1+t^2}{\sqrt{1+t^4+2t^2-4t^2}}$$

$$\frac{dx}{dt} = \frac{2(1-t^2)}{(1+t^2)^2} \times \frac{(1+t^2)}{\sqrt{1+t^4-2t^2}}$$

$$\frac{dx}{dt} = \frac{2(1-t^2)}{(1+t^2)} \times \frac{1}{\sqrt{(1-t^2)^2}}$$

$$\Rightarrow \frac{dx}{dt} = \frac{2(1-t^2)}{(1+t^2)} \times \frac{1}{(1-t^2)} \Rightarrow \frac{dx}{dt} = \frac{2}{1+t^2}$$

Now taking,  $\tan y = \frac{2t}{1-t^2}$

Differentiating both sides w.r.t  $t$ , we get

$$\frac{d}{dt}(\tan y) = \frac{d}{dt}\left(\frac{2t}{1-t^2}\right)$$

$$\sec^2 y \frac{dy}{dt} = \frac{(1-t^2) \cdot \frac{d}{dt}(2t) - 2t \cdot \frac{d}{dt}(1-t^2)}{(1-t^2)^2}$$

$$\sec^2 y \frac{dy}{dt} = \frac{(1-t^2) \cdot 2 - 2t \cdot (-2t)}{(1-t^2)^2}$$

$$\begin{aligned}
 \sec^2 y \frac{dy}{dt} &= \frac{2 - 2t^2 + 4t^2}{(1 - t^2)^2} \\
 \frac{dy}{dt} &= \frac{2 + 2t^2}{(1 - t^2)^2} \times \frac{1}{\sec^2 y} \\
 \frac{dy}{dt} &= \frac{2(1 + t^2)}{(1 - t^2)^2} \times \frac{1}{1 + \tan^2 y} \\
 \frac{dy}{dt} &= \frac{2(1 + t^2)}{(1 - t^2)^2} \times \frac{1}{1 + \left(\frac{2t}{1 - t^2}\right)^2} \\
 \frac{dy}{dt} &= \frac{2(1 + t^2)}{(1 - t^2)^2} \times \frac{1}{\frac{(1 - t^2)^2 + 4t^2}{(1 - t^2)^2}} \\
 \frac{dy}{dt} &= \frac{2(1 + t^2)}{(1 - t^2)^2} \times \frac{(1 - t^2)^2}{1 + t^2 + 2t^2 + 4t^2} \\
 \frac{dy}{dt} &= \frac{2(1 + t^2)}{(1 - t^2)^2} \times \frac{(1 - t^2)^2}{1 + t^4 + 2t^2} \\
 \Rightarrow \frac{dy}{dt} &= \frac{2(1 + t^2)}{(1 - t^2)^2} \times \frac{(1 - t^2)^2}{(1 + t^2)^2} \Rightarrow \frac{dy}{dt} = \frac{2}{1 + t^2} \\
 \therefore \frac{dy}{dt} &= \frac{dy/dt}{dx/dt} = \frac{\frac{2}{1 + t^2}}{\frac{2}{1 + t^2}} = 1
 \end{aligned}$$

Thus,  $\frac{dy}{dt} = 1$

$$x = \frac{1 + \log t}{t^2}, \quad y = \frac{3 + 2 \log t}{t}$$

48.

**Solution:**

On differentiating both the given parametric functions w.r.t.  $t$ , we have

$$\begin{aligned}
 \frac{dx}{dt} &= \frac{t^2 \cdot \frac{d}{dt}(1 + \log t) - (1 + \log t) \cdot \frac{d}{dt}(t^2)}{t^4} \\
 &= \frac{t^2 \cdot \left(\frac{1}{t}\right) - (1 + \log t) \cdot 2t}{t^4} = \frac{t - (1 + \log t) \cdot 2t}{t^4} \\
 &= \frac{t[1 - 2 - 2 \log t]}{t^4} = \frac{-(1 + 2 \log t)}{t^3}
 \end{aligned}$$

$$y = \frac{3 + 2 \log t}{t}$$

Next,

$$\frac{dy}{dt} = \frac{t \cdot \frac{d}{dt}(3 + 2 \log t) - (3 + 2 \log t) \cdot \frac{d}{dt}(t)}{t^2}$$

$$= \frac{t(2/t) - (3 + 2 \log t) \cdot 1}{t^2}$$

$$= \frac{2 - 3 - 2 \log t}{t^2} = \frac{-(1 + 2 \log t)}{t^2}$$

$$\therefore \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{\frac{-(1 + 2 \log t)}{t^2}}{\frac{t^3}{t^2}} = \frac{t^3}{t^2} = t$$

Thus,  $\frac{dy}{dx} = t$ .

**49. If  $x = e^{\cos 2t}$  and  $y = e^{\sin 2t}$ , prove that  $dy/dx = -y \log x / x \log y$ .**

**Solution:**

Given,

$$x = e^{\cos 2t} \text{ and } y = e^{\sin 2t}$$

So,  $\cos 2t = \log x$  and  $\sin 2t = \log y$

Now, differentiating both the parameter functions w.r.t  $t$ , we have

$$\begin{aligned} \frac{dx}{dt} &= e^{\cos 2t} \cdot \frac{d}{dt}(\cos 2t) = e^{\cos 2t} (-\sin 2t) \cdot \frac{d}{dt}(2t) \\ &= -e^{\cos 2t} \cdot \sin 2t \cdot 2 = -2e^{\cos 2t} \cdot \sin 2t \end{aligned}$$

$$\text{Now, } y = e^{\sin 2t}$$

$$\begin{aligned} \frac{dy}{dt} &= e^{\sin 2t} \cdot \frac{d}{dt}(\sin 2t) = e^{\sin 2t} \cdot \cos 2t \cdot \frac{d}{dt}(2t) \\ &= e^{\sin 2t} \cdot \cos 2t \cdot 2 = 2e^{\sin 2t} \cdot \cos 2t \end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2e^{\sin 2t} \cdot \cos 2t}{-2e^{\cos 2t} \cdot \sin 2t} = \frac{e^{\sin 2t} \cdot \cos 2t}{-e^{\cos 2t} \cdot \sin 2t} = \frac{y \cos 2t}{-x \sin 2t}$$

$$= \frac{y \log x}{-x \log y} \quad \left[ \begin{array}{l} \because \cos 2t = \log x \\ \sin 2t = \log y \end{array} \right]$$

$$\text{Thus, } \frac{dy}{dx} = -\frac{y \log x}{x \log y}.$$

**50. If  $x = a \sin 2t (1 + \cos 2t)$  and  $y = b \cos 2t (1 - \cos 2t)$ , show that  $\left( \frac{dy}{dx} \right)_{at t = \frac{\pi}{4}} = \frac{b}{a}$ .**

Given,

$$x = a \sin 2t (1 + \cos 2t) \text{ and } y = b \cos 2t (1 - \cos 2t)$$

Differentiating both the parametric equations w.r.t  $t$ , we have

$$\begin{aligned}
 \frac{dx}{dt} &= a \left[ \sin 2t \cdot \frac{d}{dt}(1 + \cos 2t) + (1 + \cos 2t) \cdot \frac{d}{dt} \sin 2t \right] \\
 &= a [\sin 2t \cdot (-\sin 2t) \cdot 2 + (1 + \cos 2t)(\cos 2t) \cdot 2] \\
 &= a [-2 \sin^2 2t + 2 \cos 2t + 2 \cos^2 2t] \\
 &= a [2(\cos^2 2t - \sin^2 2t) + 2 \cos 2t] \\
 &= a [2 \cos 4t + 2 \cos 2t] \quad [\because \cos 2x = \cos^2 x - \sin^2 x] \\
 &= 2a [\cos 4t + \cos 2t]
 \end{aligned}$$

Now,  $y = b \cos 2t (1 - \cos 2t)$

$$\begin{aligned}
 \frac{dy}{dt} &= b \left[ \cos 2t \cdot \frac{d}{dt}(1 - \cos 2t) + (1 - \cos 2t) \cdot \frac{d}{dt}(\cos 2t) \right] \\
 &= b [\cos 2t \cdot \sin 2t \cdot 2 + (1 - \cos 2t) \cdot (-\sin 2t) \cdot 2] \\
 &= b [2 \sin 2t \cdot \cos 2t - 2 \sin 2t + 2 \sin 2t \cos 2t] \\
 &= b [\sin 4t - 2 \sin 2t + \sin 4t] \quad [\because \sin 2x = 2 \sin x \cos x] \\
 &= b [2 \sin 4t - 2 \sin 2t] = 2b (\sin 4t - \sin 2t)
 \end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2b [\sin 4t - \sin 2t]}{2a [\cos 4t + \cos 2t]} = \frac{b}{a} \left[ \frac{\sin 4t - \sin 2t}{\cos 4t + \cos 2t} \right]$$

Putting,  $t = \frac{\pi}{4}$  we get

$$\begin{aligned}
 \left( \frac{dy}{dx} \right)_{at t = \frac{\pi}{4}} &= \frac{b}{a} \left[ \frac{\sin 4\left(\frac{\pi}{4}\right) - \sin 2\left(\frac{\pi}{4}\right)}{\cos 4\left(\frac{\pi}{4}\right) + \cos 2\left(\frac{\pi}{4}\right)} \right] = \frac{b}{a} \left[ \frac{\sin \pi - \sin \frac{\pi}{2}}{\cos \pi + \cos \frac{\pi}{2}} \right] \\
 &= \frac{b}{a} \left[ \frac{0 - 1}{-1 + 0} \right] = \frac{b}{a} \left( \frac{-1}{-1} \right) = \frac{b}{a}
 \end{aligned}$$

Thus,  $\left( \frac{dy}{dx} \right)_{at t = \frac{\pi}{4}} = \frac{b}{a}$ .

**51. If  $x = 3 \sin t - \sin 3t$ ,  $y = 3 \cos t - \cos 3t$ , find  $\frac{dy}{dx}$  at  $t = \frac{\pi}{3}$ .**

**Solution:**

Given,

$$x = 3 \sin t - \sin 3t, y = 3 \cos t - \cos 3t$$

Now, differentiating both the parametric functions w.r.t  $t$ , we have

$$\frac{dx}{dt} = 3 \cos t - \cos 3t \cdot 3 = 3(\cos t - \cos 3t)$$

$$\frac{dy}{dt} = -3 \sin t + \sin 3t \cdot 3 = 3(-\sin t + \sin 3t)$$



$$\text{So, } \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{3(-\sin t + \sin 3t)}{3(\cos t - \cos 3t)} = \frac{-\sin t + \sin 3t}{\cos t - \cos 3t}$$

$$\text{Putting, } t = \frac{\pi}{3}$$

$$\begin{aligned} \frac{dy}{dx} &= \frac{-\sin \frac{\pi}{3} + \sin 3\left(\frac{\pi}{3}\right)}{\cos \frac{\pi}{3} - \cos 3\left(\frac{\pi}{3}\right)} \\ &= \frac{-\frac{\sqrt{3}}{2} + \sin \pi}{\frac{1}{2} - \cos \pi} = \frac{-\frac{\sqrt{3}}{2} + 0}{\frac{1}{2} - (-1)} = \frac{-\frac{\sqrt{3}}{2}}{\frac{1}{2} + 1} = \frac{-\frac{\sqrt{3}}{2}}{\frac{3}{2}} = \frac{-1}{\sqrt{3}} \end{aligned}$$

$$\text{Thus, } \frac{dy}{dx} = \frac{-1}{\sqrt{3}}.$$

**52. Differentiate  $x/\sin x$  w.r.t  $\sin x$ .**

**Solution:**

$$\text{Let } y = \frac{x}{\sin x} \text{ and } z = \sin x.$$

Differentiating both the parametric functions w.r.t  $x$ ,

$$\begin{aligned} \frac{dy}{dx} &= \frac{\sin x \cdot \frac{d}{dx}(x) - x \cdot \frac{d}{dx}(\sin x)}{(\sin x)^2} \\ &= \frac{\sin x \cdot 1 - x \cdot \cos x}{\sin^2 x} = \frac{\sin x - x \cos x}{\sin^2 x} \end{aligned}$$

$$\frac{dz}{dx} = \cos x$$

$$\begin{aligned} \text{Now, } \frac{dy}{dz} &= \frac{dy/dx}{dz/dx} = \frac{\frac{\sin x - x \cos x}{\sin^2 x}}{\cos x} = \frac{\sin x - x \cos x}{\sin^2 x \cos x} \\ &= \frac{\frac{\sin x}{\sin^2 x \cos x} - \frac{x \cos x}{\sin^2 x \cos x}}{\sin^2 x \cos x} \\ &= \frac{\frac{\tan x}{\sin^2 x} - \frac{x}{\sin^2 x}}{\sin^2 x} = \frac{\tan x - x}{\sin^2 x} \end{aligned}$$

$$\text{Thus, } \frac{dy}{dz} = \frac{\tan x - x}{\sin^2 x}.$$

**53. Differentiate  $\tan^{-1} \left( \frac{\sqrt{1+x^2}-1}{x} \right)$  w.r.t  $\tan^{-1} x$  when  $x \neq 0$ .**

**Solution:**



Let  $y = \tan^{-1} \left( \frac{\sqrt{1+x^2}-1}{x} \right)$  and  $z = \tan^{-1} x$

Now, put  $x = \tan \theta$

$\therefore y = \tan^{-1} \left( \frac{\sqrt{1+\tan^2 \theta}-1}{\tan \theta} \right)$  and  $z = \tan^{-1}(\tan \theta) = \theta$ .

So,  $\tan \left( \frac{\sqrt{\sec \theta}-1}{\tan \theta} \right) = \tan^{-1} \left( \frac{\sec \theta-1}{\tan \theta} \right)$

$$\tan^{-1} \left( \frac{\frac{1}{\cos \theta}-1}{\frac{\sin \theta}{\cos \theta}} \right) = \tan^{-1} \left( \frac{1-\cos \theta}{\sin \theta} \right)$$

$$\tan^{-1} \left( \frac{2 \sin^2 \theta/2}{2 \sin \theta/2 \cos \theta/2} \right) = \tan^{-1} \left( \frac{\sin \theta/2}{\cos \theta/2} \right)$$

$$\Rightarrow y = \tan^{-1} \left( \tan \frac{\theta}{2} \right) \Rightarrow y = \frac{\theta}{2}$$

Differentiating both parametric functions w.r.t.  $\theta$

$$\frac{dy}{d\theta} = \frac{1}{2} \cdot \frac{d}{d\theta}(\theta) \quad \text{and} \quad \frac{dz}{d\theta} = \frac{d}{d\theta}(\theta)$$

$$= \frac{1}{2} \cdot 1 = \frac{1}{2} \quad \text{and} \quad \frac{dz}{d\theta} = 1$$

$$\text{Thus, } \frac{dy}{dz} = \frac{dy/d\theta}{dz/d\theta} = \frac{1/2}{1} = \frac{1}{2}.$$

**NAGEEN GROUP**  
tamso ma jyotirgamaya

# **CLICK TO DOWNLOAD NCERT EXEMPLAR SOLUTIONS**

**CLASS 12: MATHEMATICS (ALL CHAPTERS)**





# JOIN OUR WHATSAPP GROUPS

FOR FREE EDUCATIONAL  
RESOURCES

---





## JOIN SCHOOL OF EDUCATORS WHATSAPP GROUPS FOR FREE EDUCATIONAL RESOURCES

We are thrilled to introduce the School of Educators WhatsApp Group, a platform designed exclusively for educators to enhance your teaching & Learning experience and learning outcomes. Here are some of the key benefits you can expect from joining our group:

### BENEFITS OF SOE WHATSAPP GROUPS

---

- **Abundance of Content:** Members gain access to an extensive repository of educational materials tailored to their class level. This includes various formats such as PDFs, Word files, PowerPoint presentations, lesson plans, worksheets, practical tips, viva questions, reference books, smart content, curriculum details, syllabus, marking schemes, exam patterns, and blueprints. This rich assortment of resources enhances teaching and learning experiences.
- **Immediate Doubt Resolution:** The group facilitates quick clarification of doubts. Members can seek assistance by sending messages, and experts promptly respond to queries. This real-time interaction fosters a supportive learning environment where educators and students can exchange knowledge and address concerns effectively.
- **Access to Previous Years' Question Papers and Topper Answers:** The group provides access to previous years' question papers (PYQ) and exemplary answer scripts of toppers. This resource is invaluable for exam preparation, allowing individuals to familiarize themselves with the exam format, gain insights into scoring techniques, and enhance their performance in assessments.

- **Free and Unlimited Resources:** Members enjoy the benefit of accessing an array of educational resources without any cost restrictions. Whether its study materials, teaching aids, or assessment tools, the group offers an abundance of resources tailored to individual needs. This accessibility ensures that educators and students have ample support in their academic endeavors without financial constraints.
- **Instant Access to Educational Content:** SOE WhatsApp groups are a platform where teachers can access a wide range of educational content instantly. This includes study materials, notes, sample papers, reference materials, and relevant links shared by group members and moderators.
- **Timely Updates and Reminders:** SOE WhatsApp groups serve as a source of timely updates and reminders about important dates, exam schedules, syllabus changes, and academic events. Teachers can stay informed and well-prepared for upcoming assessments and activities.
- **Interactive Learning Environment:** Teachers can engage in discussions, ask questions, and seek clarifications within the group, creating an interactive learning environment. This fosters collaboration, peer learning, and knowledge sharing among group members, enhancing understanding and retention of concepts.
- **Access to Expert Guidance:** SOE WhatsApp groups are moderated by subject matter experts, teachers, or experienced educators can benefit from their guidance, expertise, and insights on various academic topics, exam strategies, and study techniques.

Join the School of Educators WhatsApp Group today and unlock a world of resources, support, and collaboration to take your teaching to new heights. To join, simply click on the group links provided below or send a message to +91-95208-77777 expressing your interest.

**Together, let's empower ourselves & Our Students and  
inspire the next generation of learners.**

**Best Regards,  
Team  
School of Educators**



# Join School of Educators WhatsApp Groups

You will get Pre- Board Papers PDF, Word file, PPT, Lesson Plan, Worksheet, practical tips and Viva questions, reference books, smart content, curriculum, syllabus, marking scheme, toppers answer scripts, revised exam pattern, revised syllabus, Blue Print etc. here . Join Your Subject / Class WhatsApp Group.

## Kindergarten to Class XII (For Teachers Only)



[Click Here to Join](#)

**Class 1**



[Click Here to Join](#)

**Class 2**



[Click Here to Join](#)

**Class 3**



[Click Here to Join](#)

**Class 4**



[Click Here to Join](#)

**Class 5**



[Click Here to Join](#)

**Class 6**



[Click Here to Join](#)

**Class 7**



[Click Here to Join](#)

**Class 8**



[Click Here to Join](#)

**Class 9**



[Click Here to Join](#)

**Class 10**



[Click Here to Join](#)

**Class 11 (Science)**



[Click Here to Join](#)

**Class 11 (Humanities)**



[Click Here to Join](#)

**Class 11 (Commerce)**



[Click Here to Join](#)

**Class 12 (Science)**



[Click Here to Join](#)

**Class 12 (Humanities)**



[Click Here to Join](#)

**Class 12 (Commerce)**



[Click Here to Join](#)

**Kindergarten**

# Subject Wise Secondary and Senior Secondary Groups (IX & X For Teachers Only)

## Secondary Groups (IX & X)



[Click Here to Join](#)

SST



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Science



[Click Here to Join](#)

English



[Click Here to Join](#)

Hindi-A



[Click Here to Join](#)

IT Code-402



[Click Here to Join](#)

Hindi-B



[Click Here to Join](#)

Artificial Intelligence

## Senior Secondary Groups (XI & XII For Teachers Only)



[Click Here to Join](#)

Physics



[Click Here to Join](#)

Chemistry



[Click Here to Join](#)

English



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Biology



[Click Here to Join](#)

Accountancy



[Click Here to Join](#)

Economics



[Click Here to Join](#)

BST



[Click Here to Join](#)

History



[Click Here to Join](#)

**Geography**



[Click Here to Join](#)

**Sociology**



[Click Here to Join](#)

**Hindi Elective**



[Click Here to Join](#)

**Hindi Core**



[Click Here to Join](#)

**Home Science**



[Click Here to Join](#)

**Sanskrit**



[Click Here to Join](#)

**Psychology**



[Click Here to Join](#)

**Political Science**



[Click Here to Join](#)

**Painting**



[Click Here to Join](#)

**Vocal Music**



[Click Here to Join](#)

**Comp. Science**



[Click Here to Join](#)

**IP**



[Click Here to Join](#)

**Physical Education**



[Click Here to Join](#)

**APP. Mathematics**



[Click Here to Join](#)

**Legal Studies**



[Click Here to Join](#)

**Entrepreneurship**



[Click Here to Join](#)

**French**



[Click Here to Join](#)

**IT**



[Click Here to Join](#)

**Artificial Intelligence**

## **Other Important Groups (For Teachers & Principal's)**



[Click Here to Join](#)

**Principal's Group**



[Click Here to Join](#)

**Teachers Jobs**



[Click Here to Join](#)

**IIT/NEET**

# Join School of Educators WhatsApp Groups

You will get Pre- Board Papers PDF, Word file, PPT, Lesson Plan, Worksheet, practical tips and Viva questions, reference books, smart content, curriculum, syllabus, marking scheme, toppers answer scripts, revised exam pattern, revised syllabus, Blue Print etc. here . Join Your Subject / Class WhatsApp Group.

## Kindergarten to Class XII (For Students Only)



[Click Here to Join](#)

**Class 1**



[Click Here to Join](#)

**Class 2**



[Click Here to Join](#)

**Class 3**



[Click Here to Join](#)

**Class 4**



[Click Here to Join](#)

**Class 5**



[Click Here to Join](#)

**Class 6**



[Click Here to Join](#)

**Class 7**



[Click Here to Join](#)

**Class 8**



[Click Here to Join](#)

**Class 9**



[Click Here to Join](#)

**Class 10**



[Click Here to Join](#)

**Class 11 (Science)**



[Click Here to Join](#)

**Class 11 (Humanities)**



[Click Here to Join](#)

**Class 11 (Commerce)**



[Click Here to Join](#)

**Class 12 (Science)**



[Click Here to Join](#)

**Class 12 (Humanities)**



[Click Here to Join](#)

**Class 12 (Commerce)**



[Click Here to Join](#)

**Artificial Intelligence  
(VI TO VIII)**

# Subject Wise Secondary and Senior Secondary Groups (IX & X For Students Only) Secondary Groups (IX & X)



[Click Here to Join](#)

SST



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Science



[Click Here to Join](#)

English



[Click Here to Join](#)

Hindi



[Click Here to Join](#)

IT Code



[Click Here to Join](#)

Artificial Intelligence

## Senior Secondary Groups (XI & XII For Students Only)



[Click Here to Join](#)

Physics



[Click Here to Join](#)

Chemistry



[Click Here to Join](#)

English



[Click Here to Join](#)

Mathematics



[Click Here to Join](#)

Biology



[Click Here to Join](#)

Accountancy



[Click Here to Join](#)

Economics



[Click Here to Join](#)

BST



[Click Here to Join](#)

History





[Click Here to Join](#)

**Geography**



[Click Here to Join](#)

**Sociology**



[Click Here to Join](#)

**Hindi Elective**



[Click Here to Join](#)

**Hindi Core**



[Click Here to Join](#)

**Home Science**



[Click Here to Join](#)

**Sanskrit**



[Click Here to Join](#)

**Psychology**



[Click Here to Join](#)

**Political Science**



[Click Here to Join](#)

**Painting**



[Click Here to Join](#)

**Music**



[Click Here to Join](#)

**Comp. Science**



[Click Here to Join](#)

**IP**



[Click Here to Join](#)

**Physical Education**



[Click Here to Join](#)

**APP. Mathematics**



[Click Here to Join](#)

**Legal Studies**



[Click Here to Join](#)

**Entrepreneurship**



[Click Here to Join](#)

**French**



[Click Here to Join](#)

**IT**



[Click Here to Join](#)

**AI**



[Click Here to Join](#)

**IIT/NEET**



[Click Here to Join](#)

**CUET**

## Groups Rules & Regulations:

**To maximize the benefits of these WhatsApp groups, follow these guidelines:**

1. Share your valuable resources with the group.
2. Help your fellow educators by answering their queries.
3. Watch and engage with shared videos in the group.
4. Distribute WhatsApp group resources among your students.
5. Encourage your colleagues to join these groups.

### **Additional notes:**

1. Avoid posting messages between 9 PM and 7 AM.
2. After sharing resources with students, consider deleting outdated data if necessary.
3. It's a NO Nuisance groups, single nuisance and you will be removed.
  - No introductions.
  - No greetings or wish messages.
  - No personal chats or messages.
  - No spam. Or voice calls
  - Share and seek learning resources only.

**Please only share and request learning resources. For assistance, contact the helpline via WhatsApp: +91-95208-77777.**

# Join Premium WhatsApp Groups Ultimate Educational Resources!!

Join our premium groups and just Rs. 1000 and gain access to all our exclusive materials for the entire academic year. Whether you're a student in Class IX, X, XI, or XII, or a teacher for these grades, Artham Resources provides the ultimate tools to enhance learning. Pay now to delve into a world of premium educational content!

[Click here for more details](#)



**Click Here to Join**

**Class 9**



**Click Here to Join**

**Class 10**



**Click Here to Join**

**Class 11**



**Click Here to Join**

**Class 12**

📢 Don't Miss Out! Elevate your academic journey with top-notch study materials and secure your path to top scores! Revolutionize your study routine and reach your academic goals with our comprehensive resources. Join now and set yourself up for success! 🇧🇩🌟

**Best Wishes,**

**Team**

**School of Educators & Artham Resources**

# SKILL MODULES BEING OFFERED IN MIDDLE SCHOOL



Artificial Intelligence



Beauty & Wellness




Design Thinking & Innovation



Financial Literacy



Handicrafts



Information Technology



Marketing/Commercial Application




Mass Media - Being Media Literate




Travel & Tourism




Coding



Data Science (Class VIII only)




Augmented Reality / Virtual Reality



Digital Citizenship




Life Cycle of Medicine & Vaccine



Things you should know about keeping Medicines at home




What to do when Doctor is not around



Humanity & Covid-19




Blue Pottery




Pottery



Block Printing




Food



Food Preservation




Baking



Herbal Heritage



Khadi




Mask Making




Mass Media



Making of a Graphic Novel




Kashmiri Embroidery



Embroidery




Rockets



Satellites



Application of Satellites




Photography



# SKILL SUBJECTS AT SECONDARY LEVEL (CLASSES IX – X)



Retail



Information Technology




Security




Automotive



Introduction To Financial Markets



Introduction To Tourism



Beauty & Wellness



Agriculture



Food Production



Front Office Operations



Banking & Insurance



Marketing & Sales



Health Care



Apparel



Multi Media



Multi Skill Foundation Course



Artificial Intelligence



Physical Activity Trainer



Data Science



Electronics & Hardware (NEW)




Foundation Skills For Sciences (Pharmaceutical & Biotechnology)(NEW)




Design Thinking & Innovation (NEW)

# SKILL SUBJECTS AT SR. SEC. LEVEL (CLASSES XI – XII)



Retail



Information Technology



Web Application



Automotive



Financial Markets Management



Tourism



Beauty & Wellness



Agriculture



Food Production



Front Office Operations



Banking



Marketing




Health Care




Insurance




Horticulture



Typography & Comp.  
Application




Geospatial Technology



Electrical Technology



Electronic Technology




Multi-Media




Taxation




Cost Accounting




Office Procedures & Practices




Shorthand (English)




Shorthand (Hindi)




Air-Conditioning & Refrigeration




Medical Diagnostics




Textile Design




Design




Salesmanship



Business Administration



Food Nutrition & Dietetics




Mass Media Studies



Library & Information Science



Fashion Studies



Applied Mathematics



Yoga



Early Childhood Care & Education




Artificial Intelligence



Data Science



Physical Activity Trainer(new)



Land Transportation Associate (NEW)





Electronics & Hardware (NEW)



Design Thinking & Innovation (NEW)





- Pdf Purchase

- Print Purchase

- Amazon Print Purchase



Scan QR Code to join Telegram Groups & Channels for more free resources.

